

Supplement of Geosci. Model Dev., 10, 1587–1605, 2017
<http://www.geosci-model-dev.net/10/1587/2017/>
doi:10.5194/gmd-10-1587-2017-supplement
© Author(s) 2017. CC Attribution 3.0 License.



Supplement of

A framework for expanding aqueous chemistry in the Community Multiscale Air Quality (CMAQ) model version 5.1

Kathleen M. Fahey et al.

Correspondence to: Kathleen M. Fahey (fahey.kathleen@epa.gov)

The copyright of individual parts of the supplement might differ from the CC-BY 3.0 licence.

Table S1. Gas phase species* that participate in phase transfer and associated gas phase diffusion and accommodation coefficients. Unless otherwise indicated, Henry’s Law coefficients are the base model’s original values (from CMAQ subroutine hlconst.F).

5

Reaction	H _{law} (M/atm)	−ΔH/R (K)	α ^a	D _g (m ² /s) x 10 ^{5b}
SO ₂ (gas) ↔ SO ₂ (aq)	1.4	2.9E+03	1.10 x 10 ⁻¹	1.28
HNO ₃ (gas) ↔ HNO ₃ (aq)	2.1E+05	8.7E+03	8.68 x 10 ⁻²	1.32
CO ₂ (gas) ↔ H ₂ CO ₃ (aq)	3.6E-02	2.2E+03	1.50 x 10 ⁻⁴	1.55
NH ₃ (gas) ↔ NH ₄ OH (aq)	6.1E+01	4.2E+03	9.10 x 10 ⁻²	2.30
H ₂ O ₂ (gas) ↔ H ₂ O ₂ (aq)	8.3E+04	7.4E+03	1.53 x 10 ⁻¹	1.46
O ₃ (gas) ↔ O ₃ (aq)	1.14E-02	2.3E+03	1.00 x 10 ⁻¹	1.48
HCOOH (gas) ↔ HCOOH (aq)	8.9E+03	6.1E+03	2.29 x 10 ⁻²	1.53
MHP (gas) ↔ MHP (aq)	3.1E+02	5.2E+03	6.76 x 10 ⁻³	1.31
PAA (gas) ↔ PAA (aq)	8.4E+02	5.3E+03	1.90 x 10 ⁻²	1.02
HCl (gas) ↔ HCl (aq)	1.9E+01	6.0E+02	1.16 x 10 ⁻¹	1.89
GLY (gas) ↔ GLY (aq)	3.6E+05		2.30 x 10 ^{-2c}	1.15 ^a
MGLY (gas) ↔ MGLY (aq)	3.2E+04 ^e		2.30 x 10 ^{-2c}	1.15 ^d
IEPOX (gas) ↔ IEPOX (aq)	2.7E+06 ^f		2.00 x 10 ^{-2f}	1.00 ^f
MAE (gas) ↔ MAE (aq)	1.2E+05 ^f		2.00 x 10 ^{-2f}	1.00 ^f
HMML (gas) ↔ HMML (aq)	1.2E+05 ^f		2.00 x 10 ^{-2f}	1.00 ^f

H₂SO₄ is instantaneously transferred to aerosol SO₄ and N₂O₅ to HNO₃ (gas) at the start of cloud processing. ^aCAPRAM 2.4 tables (http://projects.tropos.de/capram/capram_24.html). ^bHerrmann et al. (2000)*. ^cLim et al. (2005)*. ^dAssumed equal to D_g for glyoxal. ^eThe Henry’s Law coefficient for methylglyoxal is replaced with that of glyoxal when the CB05 gas phase mechanism is used (Carlton et al., 2010). ^fPye et al. (2013). *and the original sources therein

10

Table S2. Dissociation reactions and equilibrium constants. Ionic species are considered explicitly and dissociation equilibrium reactions are described as a set of forward and backward reactions. Activity coefficients are rolled into the forward and backward reaction rates.

Reaction	K _{eq,298} ^a [M]	−ΔE ₀ /R [K] [#]	k _b ^{a,d} [M ⁻¹ s ⁻¹]
SO ₂ ↔ HSO ₃ ⁻ + H ⁺	1.39x10 ⁻²	1.87x10 ³	2.0x10 ⁸
HSO ₃ ⁻ ↔ SO ₃ ²⁻ + H ⁺	6.72x10 ⁻⁸	3.55x10 ²	5.0x10 ¹⁰
HNO ₃ ↔ NO ₃ ⁻ + H ⁺	1.7x10 ¹		5.0x10 ¹⁰
NH ₄ OH ↔ NH ₄ ⁺ + OH ⁻	1.77x10 ⁻⁵	-7.10x10 ²	3.4x10 ¹⁰
H ₂ CO ₃ ↔ HCO ₃ ⁻ + H ⁺	4.3x10 ⁻⁷	-9.95x10 ²	6.4x10 ⁴
HCO ₃ ⁻ ↔ CO ₃ ²⁻ + H ⁺	4.68x10 ⁻¹¹	-1.79x10 ³	5.0x10 ¹⁰
HCOOH ↔ HCOO ⁻ + H ⁺	1.8x10 ⁻⁴ ^b	-2.00x10 ¹	5.0x10 ¹⁰
HCl ↔ Cl ⁻ + H ⁺	1.74x10 ⁶ ^b	6.90x10 ³	5.0x10 ¹⁰
H ₂ O ↔ OH ⁻ + H ⁺	1.8x10 ⁻¹⁶ ^c	-6.95x10 ³	1.4x10 ¹¹
H ₂ SO ₄ ↔ HSO ₄ ⁻ + H ⁺	1.0x10 ³ ^b		5.0x10 ¹⁰
HSO ₄ ⁻ ↔ SO ₄ ²⁻ + H ⁺	1.02x10 ⁻²	2.45x10 ³	1.0x10 ¹¹

^aWarneck and Williams (2012)*. ^bSeinfeld and Pandis(2006)*. ^cRemoved [H₂O] concentration of 55.5 M from the original equilibrium constant. ^dGraedel and Weschler (1981)*. *and the original sources therein

15

k_r = K_{eq,i,T} x k_b [#]Note that while there are some minor differences between some of the equilibrium constants used here compared to AQCHEM, box model tests indicate that these small differences only have a minimal impact on predicted concentrations

20

Table S3. Chemical kinetic reactions.

Reaction	Rate coefficient in appropriate units of M,s (298K); -E _a /R (K) [§]		Notes
HSO ₃ ⁻ + H ₂ O ₂ + H ⁺ → SO ₄ ²⁻ + 2 H ⁺	7.45E+7 / (1 + 13 [H+]); -4756.08		
SO ₂ + O ₃ → SO ₄ ²⁻ + 2 H ⁺	2.4E+4; 0.0		Q
HSO ₃ ⁻ + O ₃ → SO ₄ ²⁻ + H ⁺	3.7E+5; -5530.88		Q
SO ₃ ²⁻ + O ₃ → SO ₄ ²⁻	1.5E+9; -5280.56		Q
HSO ₃ ⁻ + MHP + H ⁺ → SO ₄ ²⁻ + 2 H ⁺	1.9E+7; -3799.5		
*HSO ₃ ⁻ + PAA + H ⁺ → SO ₄ ²⁻ + 2 H ⁺	3.6E+7; -3999.2		*
*HSO ₃ ⁻ + PAA → SO ₄ ²⁻ + H ⁺	7E+2; 0.0		
SO ₂ (+ Mn(II)) → SO ₄ ²⁻ + 2 H ⁺	7.5E+2; 0.0		#
HSO ₃ ⁻ (+ Mn(II)) → SO ₄ ²⁻ + H ⁺	7.5E+2; 0.0		#
SO ₃ ²⁻ (+ Mn(II)) → SO ₄ ²⁻	7.5E+2; 0.0		#
SO ₂ (+ Fe(III)) → SO ₄ ²⁻ + 2 H ⁺	2.6E+3; 0.0		#
HSO ₃ ⁻ (+ Fe(III)) → SO ₄ ²⁻ + H ⁺	2.6E+3; 0.0		#
SO ₃ ²⁻ (+ Fe(III)) → SO ₄ ²⁻	2.6E+3; 0.0		#
SO ₂ (+Mn(II) + Fe(III)) → SO ₄ ²⁻ + 2 H ⁺	1E+10; 0.0		#
HSO ₃ ⁻ (+ Mn(II) + Fe(III)) → SO ₄ ²⁻ + H ⁺	1E+10; 0.0		#
SO ₃ ²⁻ (+ Mn(II) + Fe(III)) → SO ₄ ²⁻	1E+10; 0.0		#
GLY + OH• → 0.04 ORGC	3E+10; 0.0		Fixed OH•
MGLY + OH• → 0.04 ORGC	3E+10; 0.0		Fixed OH•
	k ₁ (M ⁻² s ⁻¹)	k ₂ (M ⁻² s ⁻¹)	
IEPOX + H ₂ O → 2-methyltetrol	9E-04	1.3E-05	Rate = KIEPOX * Parent
IEPOX + SO ₄ ²⁻ → IEPOX_OS	2E-04	2.9E-06	Hydrocarbon *
IEPOX + NO ₃ ⁻ → IEPOX_ON [§]	2E-04	2.9E-06	Nucleophile * Q
IEPOX + 2-methyltetrol → dimers	2E-04	2.9E-06	
IEPOX + IEPOX_OS → dimers	2E-04	2.9E-06	where
IEPOX + IEPOX_ON → dimers [§]	2E-04	2.9E-06	KIEPOX = k ₁ *[H ⁺] +
MAE + H ₂ O → 2-MG	9E-04	1.3E-05	k ₂ *[HSO ₄ ⁻]
MAE + SO ₄ ²⁻ → MPAN_OS	2E-04	2.9E-06	
MAE + NO ₃ ⁻ → MPAN_ON [§]	2E-04	2.9E-06	
MAE + 2-MG → dimers [§]	2E-04	2.9E-06	H ₂ O concentration
MAE + MPAN_OS → dimers [§]	2E-04	2.9E-06	remains fixed (55.5 M)
MAE + MPAN_ON → dimers [§]	2E-04	2.9E-06	
HMML + H ₂ O → 2-MG	9E-04	1.3E-05	
HMML + SO ₄ ²⁻ → MPAN_OS	2E-04	2.9E-06	
HMML + NO ₃ ⁻ → MPAN_ON [§]	2E-04	2.9E-06	
HMML + 2-MG → dimers [§]	2E-04	2.9E-06	
HMML + MPAN_OS → dimers [§]	2E-04	2.9E-06	
HMML + MPAN_ON → dimers [§]	2E-04	2.9E-06	

[§]Rate coefficients are equal to those used in the standard version of AQCHEM. ^QAqueous diffusion correction factor applied; ^{*}included here as two reactions to accommodate the original single rate expression which included pH dependent and pH independent elements. [#]A S(VI) inhibition factor of the following form is applied to these reaction rates: (1+75[S(VI)]^{0.67})⁻¹. (Species in parentheses on the left hand side of the rate expression indicate species that affect the reaction rate calculation but whose concentrations are not consumed in the reaction.) [§]Included here but not included in the version of AQCHEM-KMTI released with CMAQv5.1

5

Table S4. CMAQ species and local AQCHEM-KMT(I) species. The “G”, “L”, “A”, and “WD” prefixes in column 1 refer to the gas-phase, aqueous-phase, Aitken aerosol, and wet deposition portion of the species respectively. Shaded cells include species specific to AQCHEM-KMTI. CMAQ aerosol concentration and wet deposition indices are of the form: [species, mode].


AQCHEM-KMT(I) local dynamic species	CMAQ species	Name and Notes
G_SO2, L_SO2, WD_SO2	GAS(LSO2), GASWDEP(LSO2)	Sulfur dioxide
G_HNO3, L_HNO3, WD_HNO3	GAS(LHNO3)/AEROSOL(LNO3, 2/3), GASWDEP(LHNO3)	Nitric acid, nitrate
G_CO2, L_H2CO3, WD_CO2	GAS(LCO2), GASWDEP(LCO2)	Carbon dioxide
G_NH3, L_NH4OH, WD_NH4OH	GAS(LNH3), AEROSOL(LNH4, 2/3), GASWDEP(LNH3)	Ammonia, ammonium
G_H2O2, L_H2O2, WD_H2O2	GAS(LH2O2), GASWDEP(LH2O2)	Hydrogen peroxide
G_O3, L_O3, WD_O3	GAS(LO3), GASWDEP(LO3)	Ozone
G_HCOOH, L_HCOOH, WD_HCOOH	GAS(LFOA), GASWDEP(LFOA)	Formic acid
G_MHP, L_MHP, WD_MHP	GAS(LMHP), GASWDEP(LMHP)	Methylhydroperoxide (CH3OOH)
G_PAA, L_PAA, WD_PAA	GAS(LPAA), GASWDEP(LPAA)	peroxyacetic acid (CH3C(O)OOH)
G_HCL, L_HCL, WD_HCL	GAS(HCL), GASWDEP(HCL)	Hydrochloric acid
G_GLY, L_GLY, WD_GLY	GAS(LGLY), GASWDEP(LGLY)	Glyoxal
G_MGLY, L_MGLY, WD_MGLY	GAS(LMGLY), GASWDEP(LMGLY)	Methylglyoxal
A_NO3AKN	AEROSOL(LNO3,1)	Aitken nitrate
A_NH4AKN	AEROSOL(LNH4,1)	Aitken ammonium
A_CLAKN	AEROSOL(LCL,1)	Aitken chloride
A_NAAKN	AEROSOL(LNA,1)	Aitken sodium
A_SO4AKN	AEROSOL(LSO4,1)	Aitken sulfate
A_PECAKN	AEROSOL(LEC,1)	Aitken elemental carbon
A_POAAKN	AEROSOL(LPOA,1)	Aitken primary OA
A_PRIAKN	AEROSOL(LPRI,1)	Aitken other primary
L_SO4MIN2, WD_H2SO4	AEROSOL(LSO4, 2/3), AERWDEP(LSO4, 2/3)	Sulfate
L_NO3MIN, WD_NO3MIN	GAS(LHNO3), AEROSOL(LNO3, 2/3), AERWDEP(LNO3, 2/3)	Nitrate
L_NH4PLUS, WD_NH4PLUS	GAS(LNH3), AEROSOL(LNH4, 2 and 3), AERWDEP(LNH4, 2/3)	Ammonium
L_CLMIN, WD_CLMIN	AEROSOL(LCL, 2/3), AERWDEP(LCL, 2/3)	Chloride
L_PRIACC, WD_PRIACC	AEROSOL(LPRI, 2), AERWDEP(LPRI, 2)	Other primary
L_PECACC, WD_PECACC	AEROSOL(LEC, 2), AERWDEP(LEC, 2)	Elemental carbon
L_ORGC, WD_ORGC	AEROSOL(LORGC, 2), AERWDEP(LORGC, 2)	Cloud SOA
L_POAACC, WD_POAACC	AEROSOL(LPOA, 2), AERWDEP(LPOA, 2)	Primary OA
L_HPLUS, WD_HPLUS	HPWDEP	H ⁺
L_OHMIN	--	OH ⁻

L_FEPLUS3, WD_FEPLUS3%	AEROSOL(LFEACC, 2), AERWDEP(LFEACC, 2), AEROSOL(LSOILC, 3), AERWDEP(LSOILC, 3), AEROSOL(LANTHC, 3), AERWDEP(LANTHC, 3)	Soluble iron ⁽³⁺⁾ ; %AQCHEM species include contributions from coarse aerosol
L_MNPLUS2, WD_MNPLUS2%	AEROSOL(LMNACC, 2), AERWDEP(LMNACC, 2), AEROSOL(LSOILC, 3), AERWDEP(LSOILC, 3), AEROSOL(LANTHC, 3), AERWDEP(LANTHC, 3)	Soluble manganese ⁽²⁺⁾ ; %AQCHEM species include contributions from coarse aerosol
L_HSO3MIN, (WD_SO2)	GAS(LSO2)	Bisulfite
L_SO3MIN2, (WD_SO2)	GAS(LSO2)	Sulfite
L_HCO3MIN, (WD_CO2)	GAS(LCO2)	Bicarbonate
L_CO3MIN2, (WD_CO2)	GAS(LCO2)	Carbonate
L_HCOOMIN, (WD_HCOOH)	GAS(LFOA)	Formate
L_HSO4MIN, (WD_H2SO4)	AEROSOL(LSO4, 2/3)	Bisulfate
L_CAPLUS2, WD_CAPLUS2%	AEROSOL(LCAACC, 2), AERWDEP(LCAACC, 2), AEROSOL(LSOILC, 3), AERWDEP(LSOILC, 3), AEROSOL(LANTHC, 3), AERWDEP(LANTHC, 3), AEROSOL(LSEASC, 3), AERWDEP(LSEASC, 3)	Calcium ⁽²⁺⁾ ; %AQCHEM species include contributions from coarse aerosol
L_MGPLUS2, WD_MGPLUS2%	AEROSOL(LMGACC, 2), AERWDEP(LMGACC, 2), AEROSOL(LSOILC, 3), AERWDEP(LSOILC, 3), AEROSOL(LANTHC, 3), AERWDEP(LANTHC, 3), AEROSOL(LSEASC, 3), AERWDEP(LSEASC, 3)	Magnesium ⁽²⁺⁾ ; %AQCHEM species include contributions from coarse aerosol
L_KPLUS, WD_KPLUS%	AEROSOL(LKACC, 2), AERWDEP(LKACC, 2), AEROSOL(LSOILC, 3), AERWDEP(LSOILC, 3), AEROSOL(LANTHC, 3), AERWDEP(LANTHC, 3), AEROSOL(LSEASC, 3), AERWDEP(LSEASC, 3)	Potassium ⁽⁺⁾ ; %AQCHEM species include contributions from coarse aerosol
L_NAPLUS, WD_NAPLUS%	AEROSOL(LNA, 2), AERWDEP(LNA, 2), AEROSOL(LSOILC, 3), AERWDEP(LSOILC, 3), AEROSOL(LANTHC, 3), AERWDEP(LANTHC, 3), AEROSOL(LSEASC, 3), AERWDEP(LSEASC, 3)	Sodium ⁽⁺⁾ ; %AQCHEM species include contributions from coarse aerosol
G_IEPOX, L_IEPOX, WD_IEPOX	GAS(LIEPOX), GASWDEP(LIEPOX)	Isoprene epoxydiols

G_IMAE, L_IMAE, WD_IMAE	GAS(LIMAE), GASWDEP(LIMAE)	Methacrylic acid epoxide
G_IHMML, L_IHMML, WD_IHMML	GAS(LIHMML), GASWDEP(LIHMML)	Hydroxymethylmethyl- α -lactone
L_IETET, WD_IETET	AEROSOL(LIETET, 2), AERWDEP(LIETET, 2)	2-methyltetrols
L_IEOS, WD_IEOS	AEROSOL(LIEOS, 2), AERWDEP(LIEOS, 2)	IEPOX-derived organosulfate
L_IEON, WD_IEON**	AEROSOL(LIEON, 2), AERWDEP(LIEON, 2)	IEPOX-derived organonitrate. **not included in CMAQv5.1 but used here
L_DIMER, WD_DIMER	AEROSOL(LDIMER, 2), AERWDEP(LDIMER, 2)	dimers
L_IMGA, WD_IMGA	AEROSOL(LIMGA, 2), AERWDEP(LIMGA, 2)	2-methylglyceric acid
L_IMOS, WD_IMOS	AEROSOL(LIMOS, 2), AERWDEP(LIMOS, 2)	MPAN-derived organosulfate
L_IMON, WD_IMON**	AEROSOL(LIMON, 2), AERWDEP(LIMON, 2)	MPAN-derived organonitrate. ** not included in CMAQv5.1 but used here
G_OH, L_OH*	GAS(LHO)	Hydroxyl radical. *Mapped in from CMAQ but kept constant during/after cloud processing

5 Table S5. Box model test scenarios. Initial aerosol species concentrations are held constant for the variable scenarios and are associated with an initial droplet pH of ~5. Default gas concentrations were set at 1 ppb and modal aerosol species concentrations were set to 0.1 $\mu\text{g}/\text{m}^3$, with the following exceptions: $\text{O}_3 = 50$ ppb, $\text{CO}_2 = 340$ ppm, $\text{H}_2\text{SO}_4 = 0.1$ ppb, $\text{N}_2\text{O}_5 = 0.1$ ppb, $\text{SO}_4^{2-} = 4.2$ $\mu\text{g}/\text{m}^3$, Primary inorganic (inert) = 2.1 $\mu\text{g}/\text{m}^3$, Primary OA = 1.1 $\mu\text{g}/\text{m}^3$, SOA = 2.1 $\mu\text{g}/\text{m}^3$, EC = 1.1 $\mu\text{g}/\text{m}^3$, $\text{Cl}^- = 2.3$ $\mu\text{g}/\text{m}^3$, $\text{NH}_4^+ = 2.2$ $\mu\text{g}/\text{m}^3$, $\text{NO}_3^- = 4.1$ $\mu\text{g}/\text{m}^3$, Na, accumulation mode Fe, Mn, Ca = 0.3 $\mu\text{g}/\text{m}^3$, coarse soil and anthropogenic coarse = 2 $\mu\text{g}/\text{m}^3$, coarse sea salt cations = 1 $\mu\text{g}/\text{m}^3$, Pressure = 1 atm.

Variable input	Values
Liquid Water content (g/m^3)	0.05, 0.1, 0.2, 0.3
Precipitation rate (mm/hr)	0, 2
Temperature (K)	270, 283
Daylight (binary)	0, 1
Integration time (s)	60, 300, 720
SO_2 (ppb)	1, 5, 10
H_2O_2 (ppb)	0.1, 1, 5
(M)GLY (ppb)	0.5, 2
OH (ppb)	10^{-3} , 10^{-5} , 10^{-7}
NH_3 (ppb)	1, 5
HNO_3 (ppb)	1, 5



5 **Figure S1: Cloud SOA (ORGC) predicted by AQCHEM-KMT at 5 (blue) and 30 μm (red) droplet diameters versus standard AQCHEM with a forced small time step. Compared to Figure 2 where AQCHEM is run with its standard time-stepping procedure, it appears that the differences between the models for ORGC are driven by both mass transfer limitations as well as the time-stepping procedure (which in standard AQCHEM is based on sulfate production only).**




Figure S2: Predicted SO₄²⁻ concentrations for AQCHEM-KMT for initial pH of 7 versus 3.

References:

- 5 Carlton, A.G., Bhawe, P.V., Napelenok, S.L., Edney, E.O., Sarwar, G., Pinder, R.W., Pouliot, G.A., and Houyoux, M.: Model representation of secondary organic aerosol in CMAQv4.7, *Environ. Sci. Technol.*, 44, 8553-8560, doi: 10.1021/es100636q, 2010.

Graedel, T.E. and Weschler, C.J.: Chemistry within aqueous atmospheric aerosols and raindrops, *Reviews of Geophysics*, 19(4), 505-539, doi:10.1029/RG019i004p00505, 1981.

- 10 Herrmann, H., Ervens, B., Jacobi, H.-W., Wolke, R., Nowacki, P., and Zellner, R.: A chemical aqueous phase radical mechanism for tropospheric chemistry, *J. Atmos. Chem.*, 36, 231–284, doi:10.1023/A:1006318622743, 2000.

Lim, H., Carlton, A.G., and Turpin, B.J.: Isoprene forms secondary organic aerosol through cloud processing: model simulations, *Environ. Sci. Technol.*, 39, 4441-4446, doi:10.1021/es048039h, 2005.

- 15 Pye, H.O.T., Pinder, R.W., Piletic, P.R., Xie, Y., Capps, S. L., Lin, Y., Surratt, J.D., Zhang, Z., Gold, A., Luecken, D.J., Hutzell, W.T., Jaoui, M., Offenberg, J.H., Kleindienst, T.E., Lewandowski M., and Edney, E.O.: Epoxide pathways improve model predictions of isoprene markers and reveal key role of acidity in aerosol formation, *Environ. Sci. Technol.*, 47, 11056-11064, doi: 10.1021/es402106h, 2013.

Seinfeld, J. H. and Pandis, S. N.: *Atmospheric Chemistry and Physics: From Air Pollution to Climate Change*, 2nd Edn., John Wiley and Sons, Inc., 2006.

- 20 Warneck, P. and Williams, J.: *The Atmospheric Chemist's Companion: Numerical Data for Use in the Atmospheric Sciences*, Springer, 2012.